

2019 QCDR MEASURE

CMS Measure ID IROMS13 **National Quality Forum (NQF) number** NA

Name or title of measure Failure to Progress (FTP): Proportion of patients not achieving a Minimal Clinically Important Difference (MCID) to indicate functional improvement in rehabilitation of patients with hip, leg or ankle injuries using the validated Lower Extremity Function Scale (LEFS) score, or equivalent instrument which has undergone peer reviewed published validation and demonstrates a peer reviewed published MCID

Measure Description The proportion of patients failing to achieve an MCID of nine (9) points or more improvement in the LEFS change score for patients with hip, leg, or ankle injuries treated during the observation period will be reported.

Additionally, a risk-adjusted MCID proportional difference will be determined by calculating the difference between the risk model predicted and observed MCID proportion will be reported for each physical therapist or physical therapy group. The risk adjustment will be calculated using a logistic regression model using: baseline LEFS score, baseline pain score, age, sex, payer, and symptom duration (time from surgery or injury to baseline physical therapy visit).

These measures will serve as a physical or occupational therapy performance measure at the eligible physical or occupational therapist or physical or occupational therapy group level.

National Quality Strategy (NQS) domain Effective Clinical Care

Measure type Patient Reported Outcome (PRO)

Meaningful Measure Area Patient Reported Functional Outcomes

High priority status Yes

Denominator The total number of patients with hip, leg, or ankle injury evaluated and treated by a physical therapist (PT) or Occupational Therapist (OT), or PT or OT Group.

Numerator The total number of patients with hip, leg, or ankle injuries to not achieve an MCID in their LEFS change score (MCID \geq 9) from their initial visit to their final visit in PT/OT practice or PT/OT group during the observation window.

Denominator exclusions Exclude patients who did not complete 2 or more surveys. Patients will be excluded if they are non-English speaking and translation services are not available, if they are unable to read or have a mental impairment that compromises their understanding.

2019 QCDR MEASURE

Denominator exceptions PT/OT can use their clinical judgement to exclude patients who are extremely medically complex, who in their experience are likely to make poor clinical progress.

Is this measure risk adjusted Yes

Number of performance rates required for measures 6 (2nd rate is reported)

- 1) Overall proportion of patients achieving an MCID in LEFS change score will be reported.
- 2) A risk-adjusted MCID proportional difference will be reported where the difference between the risk model predicted and observed MCID (measured via LEFS) proportion will be reported.

For operative (surgical) patients:

- 3) The proportion of patients not achieving an MCID in LEFS change score will be reported.
- 4) A risk-adjusted MCID proportional difference will be reported where the difference between the risk model predicted and observed MCID (measured via LEFS) proportion will be reported.

For non-operative (non-surgical) patients:

- 5) The proportion of patients not achieving an MCID in LEFS change score will be reported.
- 6) A risk-adjusted MCID proportional difference will be reported where the difference between the risk model predicted and observed MCID (measured via LEFS) proportion will be reported.

Traditional vs. inverse measure Inverse

Proportional, continuous variable, outcome, and ratio measure indicator Proportional measure